

THE MAGIC EXTRAGALACTIC SKY

BARBARA DE LOTTO (on behalf of the MAGIC Collaboration)

*University of Udine and INFN, Sezione di Trieste
Italy*

The MAGIC telescope, with its 17-m diameter mirror, is currently the largest single-dish Imaging Air Cherenkov Telescope. It is located on the Canary Island of La Palma, at an altitude of 2200 m above sea level, and is operating since 2004. The accessible energy range is in the very high energy (VHE, $E > 100$ GeV) γ -ray domain, and roughly 40% of the duty cycle is devoted to observation of extragalactic sources. Due to the lowest energy threshold (25 GeV), it can observe the deepest universe, and it is thus well suited for extragalactic observations. The strategies of extragalactic observations by MAGIC are manifold: long time monitoring of known TeV blazars, detailed study of blazars during flare states, multiwavelength campaigns on most promising targets, and search for new VHE γ -ray emitters. In this talk, highlights of observations of extragalactic objects will be reviewed.

1 Introduction

One of the major goals of ground-based γ -ray astrophysics is the study of VHE γ -ray emission from active galactic nuclei (AGN). Except for the radio galaxies M87 and Centaurus A (and possibly 3C66B), and the flat-spectrum radio quasar 3C279, all the currently known VHE γ -ray emitters in the extragalactic sky are BL Lac objects. The sensitivity of the current Imaging Atmospheric Cherenkov Telescopes (IACT) has recently enabled detailed studies of these sources in the VHE γ -rays domain, providing information for advances in understanding the origin of the VHE γ -rays, as well as powerful tools for fundamental physics studies¹.

The IACT technique² uses the atmosphere as a calorimeter to detect the extensive air shower produced after the interaction of a VHE γ -ray. The charged particles (mainly electrons and positrons) in the air shower produce Cherenkov light that can be easily detected in the ground with photomultipliers. A Cherenkov telescope uses a large reflector area to concentrate as much as possible of these photons and focus them to a camera where an image of the atmospheric cascade is formed. By analysing this image it is possible to reconstruct the incoming direction and the energy of the γ -ray. The analysis of the images is also used to reject the much higher background of cosmic rays initiated showers.

In this paper selected results on extragalactic observations with MAGIC are presented.

2 The MAGIC telescope

MAGIC^{3,4}, located on the Canary Island of La Palma (2200 m a.s.l.), is currently the largest (17-m diameter) single-dish IACT. Due to its large collection area and uniquely designed camera, MAGIC has reached the lowest energy threshold (trigger threshold 50–60 GeV at small zenith

angles, new trigger for pulsar observations ~ 25 GeV⁵⁾ for γ -ray emission among the existing terrestrial γ -ray telescopes.

MAGIC has a sensitivity of $\sim 1.6\%$ of the Crab Nebula flux in 50 observing hours. Its energy resolution is about 30% above 100 GeV and about 25% from 200 GeV onwards. The angular resolution is 0.1 deg. The MAGIC standard analysis chain is described, e.g., in Albert et al.⁶. Observations during moderate moonshine enable a substantially extended duty cycle, which is particularly important for blazar observations. Parallel optical R -band observations are performed by the Tuorla Blazar Monitoring Program with its KVA 35-cm telescope.

A second MAGIC telescope is being commissioned⁷, which is improving the sensitivity to $\sim 0.8\%$ of Crab in 50 hours.

3 The propagation and absorption of γ -rays

While travelling long distances without deviations in the fields, VHE γ -rays suffer the absorption losses due to the interaction with the low energy photons from the extragalactic background light (EBL), limiting the distance to the source that could be detected. The standard process is $\gamma_{VHE}\gamma_{EBL} \rightarrow e^+e^-$ pair production. The corresponding cross section⁸ reaches its maximum, $\sigma_{\gamma\gamma}^{\max} \simeq 1.70 \cdot 10^{-25} \text{ cm}^2$, assuming head-on collisions, when the background photon energy is $\epsilon(E) \simeq (0.5 \text{ TeV}/E) \text{ eV}$, E being the energy of the hard (incident) photon. This shows that in the energy interval explored by the IACTs, $50 \text{ GeV} < E < 100 \text{ TeV}$, the resulting opacity is dominated by the interaction with infrared/optical/ultraviolet diffuse background photons (EBL), with $0.005 \text{ eV} < \epsilon < 10 \text{ eV}$, corresponding to the wavelength range $0.125 \mu\text{m} < \lambda < 250 \mu\text{m}$.

Based on synthetic models of the evolving stellar populations in galaxies as well as on deep galaxy counts (see, for a review,⁹), several estimates of the spectral energy distribution (SED) of the EBL have been proposed, leading to different values for the transparency of the universe to $50 \text{ GeV} < E < 100 \text{ TeV}$ photons¹⁰; the resulting uncertainties are large.

Because of the absorption produced by the EBL, the observed photon spectrum $\Phi_{\text{obs}}(E_0, z)$ is related to the emitted one $\Phi_{\text{em}}(E(z))$ by

$$\Phi_{\text{obs}}(E_0, z) = e^{-\tau_{\gamma}(E_0, z)} \Phi_{\text{em}}(E_0(1+z)) , \quad (1)$$

where E_0 is the observed energy, z the source redshift and $\tau_{\gamma}(E_0, z)$ is the optical depth¹¹.

The energy dependence of τ leads to appreciable modifications of the observed source spectrum (with respect to the spectrum at emission) even for small differences in τ , due to the exponential dependence described in Eq. (1). Since the optical depth (and consequently the absorption coefficient) increases with energy, the observed flux results steeper than the emitted one. The *horizon* (e.g. Ref. ^{12,13}) for a photon of energy E is defined as the distance corresponding to the redshift z for which $\tau(E, z) = 1$, which gives an attenuation by a factor $1/e$ (see Fig. 1). MAGIC has the lowest energy threshold, and thus is currently the best suited telescope to look farther away.

4 Multi-Wavelength Campaigns

Coordinated simultaneous multi-wavelength observations, yielding spectral energy distributions (SED) spanning over 15 decades in energy, have been recently conducted, and turn out to be essential for a deeper understanding of blazars. MAGIC participated in a number of multiwavelength-campaigns on known northern-hemisphere blazars, which involved the X-ray instruments *Suzaku* and *Swift*, the γ -ray telescopes H.E.S.S., MAGIC and VERITAS, and other optical and radio telescopes.




Figure 1: Gamma-ray horizon compared with the lower energy limit of the MAGIC and H.E.S.S. Cherenkov telescopes; the curves of the photon energy versus horizon are computed for different background evolution models by Blanch & Martinez in Ref. ¹².

- Mkn 421 was detected in two campaigns during outbursts in 2006 and 2008; the coordinated effort allowed for truly simultaneous data from optical to TeV energies, and studies of correlations between the different energy bands ^{14,15}.
- The VHE emission of PG 1553+113 showed no variability during the first multi-wavelength campaign on this blazar in July 2006 ^{16,17}; it was observed simultaneously for the first time together with AGILE during 2008 ¹⁸.
- 1ES 1959+650 showed VHE data among the lowest flux state observed from this object, while at the same time a relatively high optical and X-ray flux (both Swift/Suzaku) was found ¹⁹. The SED could be modeled assuming a one zone SSC model, using parameters similar to the ones needed for the SED measured in 2002.
- Also campaigns on 1ES 1218+304 and 1H 1426+428 have been carried out, during both of which significant X-ray variability has been observed. The VHE data are being analyzed.

Further campaigns have been and will be organized in the future.

5 Strong Flaring of Messier 87 in February 2008

M87 is the first non-blazar radio galaxy known to emit VHE γ -rays, and one of the best-studied extragalactic black-hole systems. To enable long-term studies and assess the variability timescales and the location of the VHE emission in M87, the H.E.S.S., MAGIC and VERITAS collaborations established a regular, shared monitoring of M87 and agreed on mutual alerts in case of a significant detection. During the MAGIC observations, a strong signal of 8σ significance was found on 2008 February 1st, triggering the other IACTs as well as *Swift* observations. The analysis revealed a variable (significance: 5.6σ) night-to-night γ -ray flux above 350 GeV, while no variability was found in the 150–350 GeV range ²¹. The $E > 730$ GeV short-time variability of M87 reported by ²⁰ has been confirmed. This fastest variability Δt observed so far in TeV γ -rays in M87 is on the order of or even below one day, suggesting the core of M87 as the origin of the TeV γ -rays. M87 is the first radio galaxy that shows evidence for a connection between simultaneously and well sampled radio and VHE flux variations, opening a new avenue for the study of AGN accretion and jet formation ²².

6 Blazars Detected during Optical Outbursts

MAGIC has been performing target of opportunity observations upon high optical states of known or potential VHE γ -ray emitting extragalactic sources. Up to now, this strategy has been proven very successful, with the detection of Mkn 180²³, 1ES 1011+496²⁴, and recently S5 0716+71²⁵ (paper in preparation).

In April 2008, KVA observed a high optical state of the blazar S5 0716+71, triggering MAGIC observation, which resulted in a detection of a strong 6.8σ signal, corresponding to a flux of $F_{>400\text{GeV}} \approx 10^{-11} \text{ cm}^{-2} \text{ s}^{-1}$. The MAGIC observation time was 2.6 h. The source was also in a high X-ray state²⁶.

The determination of the before-unknown redshifts of 1ES 1011+496 ($z = 0.21$)²⁴ and S5 0716+71 ($z = 0.31$)²⁷ makes these objects the third-most and second-most distant TeV blazars after 3C 279, respectively.

7 The region of 3C66A/B

The MAGIC telescope observed the region around the distant blazar 3C 66A for 54.2 h in August–December 2007. The observations resulted in the discovery of a γ -ray source centered at celestial coordinates R.A. = $2^{\text{h}}23^{\text{m}}12^{\text{s}}$ and decl. = $43^{\circ}0'.7$ (MAGIC J0223+430), coinciding with the nearby radio galaxy 3C 66B²⁸. The energy spectrum of MAGIC J0223+430 follows a power law with a normalization of $(1.7 \pm 0.3_{\text{stat}} \pm 0.6_{\text{syst}}) \times 10^{-11} \text{ TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$ at 300 GeV and a photon index $\Gamma = -3.10 \pm 0.31_{\text{stat}} \pm 0.2_{\text{syst}}$. A possible association of the excess with the blazar 3C 66A and nearby radiogalaxy 3C 66B is discussed in these proceedings²⁹.

8 Detection of the flat-spectrum radio quasar 3C 279

Observations of 3C 279, the brightest EGRET AGN³⁰, during the WEBT multi-wavelength campaign³¹ revealed a 5.77σ post-trial detection on 2006 February 23rd supported by a marginal signal on the preceding night³². The overall probability for a zero-flux lightcurve can be rejected on the 5.04σ level. Simultaneous optical *R*-band observations by the Tuorla Observatory Blazar Monitoring Program revealed that during the MAGIC observations the γ -ray source was in a generally high optical state, a factor of 2 above the long-term baseline flux, but with no indication of short time-scale variability at visible wavelengths. The observed VHE spectrum can be described by a power law with a differential photon spectral index of $\alpha = 4.1 \pm 0.7_{\text{stat}} \pm 0.2_{\text{syst}}$ between 75 and 500 GeV (Fig. 2). The measured integrated flux above 100 GeV on February 23rd is $(5.15 \pm 0.82_{\text{stat}} \pm 1.5_{\text{syst}}) \times 10^{-10} \text{ photons cm}^{-2} \text{ s}^{-1}$.

This detection extends the test on the transparency of the universe up to $z = 0.536$; the γ -ray horizon together with the IACT measurements is shown in Fig. 3 from³².

VHE observations of such distant sources were until recently impossible due to the expected strong attenuation of γ rays by the EBL, which influences the observed spectrum and flux, resulting in an exponential decrease with energy and a cutoff in the γ -ray spectrum. The reconstructed intrinsic spectrum is difficult to reconcile with models predicting high EBL densities, while low-level models, e.g. ¹⁰, are still viable. Assuming a maximum intrinsic photon index of $\alpha^* = 1.5$, an upper EBL limit is inferred, leaving a small allowed region for the EBL.

In Fig. 4 the observed values of the spectral indexes of the blazars detected so far in VHE band are shown, together with the prediction (light grey area) of the standard scenario. The recent findings suggest a higher transparency of the universe to VHE photons than expected from current models of the EBL, and could be interpreted in terms of more exotic scenarios³⁴.




Figure 2: Spectrum of 3C 279 measured by MAGIC. The grey area includes the combined statistical (1σ) and systematic errors, and underlines the marginal significance of detections at high energy. The dotted line shows compatibility of the measured spectrum with a power law of photon index $\alpha = 4.1$. The blue and red triangles are measurements corrected on the basis of the two models for the EBL density.




Figure 3: The γ -ray horizon. The redshift region over which it can be constrained by observations has been extended by MAGIC up to $z=0.536$.




Figure 4: Observed values of the spectral indexes of all the blazars detected so far in VHE band as a function of the redshift; the grey band represents the prediction for different EBL models.

9 The July-2005 Flares of Mkn 501

Mkn 501 ($z = 0.034$) is known to be a strong and variable VHE γ -ray emitter. MAGIC observed Mkn 501 for 24 nights during six weeks in summer 2005. In two of these (one with moon present), the recorded flux exceeded four times the Crab-nebula flux, and revealed rapid flux changes with doubling times as short as 3 minutes or less. For the first time, short (≈ 20 min) VHE γ -ray flares with a resolved time structure could be used for detailed studies of particle acceleration and cooling timescales. In addition, a time delay between different energy bins could be investigated, and gave some hints of a delay of the higher energy photons³⁵.

An energy-dependent speed of photons in vacuum is expected as a generic signature in some approaches to Quantum Gravity (QG) theories, where Lorentz invariance violation is a manifestation of the foamy structure of space-time at short distances. It could be reflected in modifications of the propagation of energetic particles, i.e. dispersive effects due to a non-trivial refractive index induced by the fluctuations in the space-time foam³⁸. The dependence of the speed of light on the energy E of the photon can be parameterized as

$$c' = c \left[1 \pm \left(\frac{E}{E_{S1}} \right) \pm \left(\frac{E}{E_{S2}} \right)^2 \pm \dots \right]. \quad (2)$$

The energy scales E_{S1} , E_{S2} are usually expressed in units of the Planck mass, $M_P \equiv 1.22 \times 10^{19}$ GeV/c². If the linear term dominates, Eq. (2) reduces to

$$c' = c \left[1 \pm \left(\frac{E}{E_{S1}} \right) \right]. \quad (3)$$

A favored way to search for such a dispersion relation is to compare the arrival times of photons of different energies arriving on Earth from pulses of distant astrophysical sources (see³⁹ for a review).

The reanalysis of the Mkn 501 data in³⁶ resulted in a much-improved estimate of the time-energy relation. At a zero-delay probability of $P = 0.026$, a marginal time delay of $\tau_l = (0.030 \pm 0.012)$ s GeV⁻¹ towards higher energies was found using two independent analyses, both exploiting the full statistical power of the dataset (see³⁷ for details).

Since it is not possible to exclude that this delay is due to some energy-dependent effect at the source, because the emission mechanisms are not currently understood, a lower limit of




Figure 5: Skymap of extragalactic VHE γ -ray sources together with the MAGIC field of view.

$E_{S1} > 0.21 \times 10^{18}$ GeV (95% c.l.) can be established. However, if the emission mechanism at the source were understood and the observed delays were mainly due to propagation, this number could turn into a real measurement of E_{S1} .

This pioneering study demonstrates clearly the potential scientific value of an analysis of multiple flares from different sources.

10 Conclusions

After almost 4 observation cycles, MAGIC observations of the extragalactic TeV γ -ray sources contributed to many physics insights, confirming the rich potential of VHE γ -ray astrophysics. Among the currently detected 27 VHE γ -ray emitters, MAGIC has discovered 8 new sources, and detected and studied 5 known ones.

In Fig. 5 the skymap of the detected sources, together with the MAGIC field of view⁴⁰, is shown (see this reference also for an up-to-date list).

Important contributions to the understanding of active galactic nuclei have been given, allowing both to infer the intrinsic properties of the sources and to probe the nature of photon propagation through cosmic distances.

Acknowledgments

We thank the Instituto de Astrofísica de Canarias for the excellent working conditions at the Observatorio del Roque de los Muchachos in La Palma. The support of the German BMBF and MPG, the Italian INFN, the Spanish MCINN, the ETH Research Grant TH 34/043, the Polish MNiSzW Grant N N203 390834, and the YIP of the Helmholtz Gemeinschaft is gratefully acknowledged.

References

1. A. De Angelis, O. Mansutti, M. Persic, *Nuovo Cimento* **31**, 187 (2008).
2. T.C. Weekes, Proc. Int. Workshop on "Energy Budget in the High Energy Universe", Kashiwa, Japan, Feb. 22-24, (2006).
3. C. Baixeras et al., *Nucl. Instrum. Methods A* **518**, 188 (2004).
4. J. Cortina et al., Proc. 29th ICRC, 4 (2005).
5. E. Aliu et al., *Science* **322**, 1221 (2008).

6. J. Albert et al., *Astrophys. J.* **674**, 1037 (2008).
7. M. Teshima, these Proceedings.
8. W. Heitler, *The Quantum Theory of Radiation* (Oxford: Oxford University Press) (1960).
9. M.G. Hauser and E. Dwek, *ARA&A*, **39**, 249 (2001).
10. F.W. Stecker and S.T. Scully, *Astrophys. J.* **652**, L9 (2006).
T.M. Kneiske, K. Mannheim, D.H. Hartmann, *Astron. Astrophys.* **386**, 1 (2002).
T.M.Kneiske, T. Bretz, K. Mannheim, D.H. Hartmann, *Astron. Astrophys.* **413**, 807 (2004).
J.R. Primack, J.S. Bullock, R.S. Somerville, *High Energy Gamma-Ray Astronomy*, AIP Conference Series, 745, 23 (2005).
F.A. Aharonian et al., *Astron. Astrophys.* **448**, L19 (2006).
T. Stanev, D. De Marco, M.A. Malkan, F.W. Stecker, *Phys. Rev. D* **73**, 04302-2 (2006).
D. Mazin and F. Goebel, *Astrophys. J.* **655**, L13 (2007).
A. Franceschini, G. Rodighiero, M. Vaccari, *Astron. Astrophys.* **487**, 837 (2008).
M. Raue and D. Mazin, *Int. Journ. of Mod. Phys. D* **9**, 17 (2008).
11. G.G. Fazio and F.W. Stecker, *Nature* **226**, 135 (1970).
12. O. Blanch and M. Martinez, *Astrophys. J.* **23**, 598 (2005).
13. K. Mannheim, *Rev. Mod. Astron.* **12**, 167 (1999).
14. I. Donnarumma et al., *Astrophys. J.* **691**, L13 (2009).
15. V.A. Acciari et al., paper in preparation.
16. A. Reimer et al., *Astrophys. J.* **682**, 775 (2008).
17. J. Albert et al., *Astron. Astrophys.* **RN493**, (2009).
18. paper in preparation.
19. G. Tagliaferri et al., *Astrophys. J.* **679**, 1029 (2008).
20. F. Aharonian et al., *Science* **314**, 1424 (2006).
21. J. Albert et al., *Astrophys. J.* **685**, L23 (2008).
22. V.A. Acciari et al., paper in preparation.
23. J. Albert et al., *Astrophys. J.* **648**, L105 (2006).
24. J. Albert et al., *Astrophys. J.* **667**, L21 (2007).
25. M. Teshima, *Astronomer's Telegram* 1500 (2008).
26. P. Giommi et al., *Astronomer's Telegram* 1495 (2008).
27. K. Nilsson et al., *Astron. Astrophys.* **487**, L29 (2008).
28. E. Aliu et al., *Astrophys. J.* **692**, L29 (2009).
29. D. Mazin, these Proceedings.
30. A. E. Wehrle et al., *Astrophys. J.* **497**, 178 (1998).
31. M. Böttcher et al., *Astrophys. J.* **670**, 968 (2008).
32. J. Albert et al., *Science* **320**, 1752 (2008).
33. F. Aharonian et al., *Nature* **440**, 1018 (2006).
34. A. De Angelis, O. Mansutti, M. Persic, M. Roncadelli, *Mon. Not. R. Astron.* (2009).
35. J. Albert et al., *Astrophys. J.* **669**, 862 (2007).
36. J. Albert et al., *Phys. Lett. B* **668**, 253 (2008).
37. M. Martinez, M. Errando, preprint: arXiv:0803.2120 [astro-ph], submitted.
38. G. Amelino-Camelia, J. Ellis, N.E. Mavromatos, D.V. Nanopoulos and S. Sakar, *Nature* **393**, 763 (1998).
39. D. Mattingly, *Living Rev. Rel.* **8**, 5 (2005).
40. R. M. Wagner, <http://www.mpp.mpg.de/~rwagner/sources/>.